

=> fil hcaplus

FILE 'HCAPLUS' ENTERED AT 16:36:02 ON 29 APR 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 29 Apr 2005 VOL 142 ISS 19

FILE LAST UPDATED: 28 Apr 2005 (20050428/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.


This file contains CAS Registry Numbers for easy and accurate substance identification.

=>

=>

=> d stat que

```
L39      33 SEA FILE=HCAPLUS ABB=ON  PLU=ON  ("GARZON A"/AU OR "GARZON A
          E"/AU OR "GARZON AARON"/AU)
L40      443 SEA FILE=HCAPLUS ABB=ON  PLU=ON  ("FINK G"/AU OR "FINK G A"/AU
          OR "FINK G B"/AU OR "FINK G C"/AU OR "FINK G D"/AU OR "FINK G
          E"/AU OR "FINK G J"/AU OR "FINK G R"/AU) OR ("FINK GEORG"/AU
          OR "FINK GEORGE"/AU OR "FINK GEORGE A"/AU)
L41      6 SEA FILE=HCAPLUS ABB=ON  PLU=ON  L39 AND L40
L42      TRANSFER PLU=ON  L41 1-6 RN :      199 TERMS
L43      199 SEA FILE=REGISTRY ABB=ON  PLU=ON  L42
L47      STR
```



NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 14

STEREO ATTRIBUTES: NONE

L49 38 SEA FILE=REGISTRY SUB=L43 SSS FUL L47
 L50 4501 SEA FILE=HCAPLUS ABB=ON PLU=ON L49
 L51 652 SEA FILE=REGISTRY ABB=ON PLU=ON CANNABI?
 L52 15590 SEA FILE=HCAPLUS ABB=ON PLU=ON L51 OR ?CANNABI?
 L53 4475 SEA FILE=HCAPLUS ABB=ON PLU=ON L50 AND L52
 L54 182 SEA FILE=HCAPLUS ABB=ON PLU=ON L53 AND ?PSYCHOTRO?
 L55 134 SEA FILE=HCAPLUS ABB=ON PLU=ON L54 AND PD=<JULY 28, 2000
 L56 32 SEA FILE=HCAPLUS ABB=ON PLU=ON (NONPSYCHO? OR NON(W).PSYCHOTRO
 ?) AND L55

=>

=>

=> d ibib abs hitstr l56 1-32

L56 ANSWER 1 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 2001:13489 HCAPLUS
 DOCUMENT NUMBER: 135:70407
 TITLE: Dexanabinol Pharmos
 AUTHOR(S): Pop, Emil
 CORPORATE SOURCE: Alchem Laboratories Corporation, Alachua, FL, 32615,
 USA
 SOURCE: Current Opinion in Investigational Drugs (PharmaPress
 Ltd.) (2000), 1(4), 494-503
 CODEN: COIDAZ
 PUBLISHER: PharmaPress Ltd.
 DOCUMENT TYPE: Journal; General Review
 LANGUAGE: English

AB A review, with 106 refs. Dexanabinol is a **non-
 psychotropic cannabinoid** NMDA receptor antagonist under
 development by Pharmos Corp for the potential treatment of cerebral
 ischemia, glaucoma, Alzheimer's disease, cardiac failure, head injury and
 multiple sclerosis (MS); it is in phase III trials for traumatic brain
 injury (TBI). Dexanabinol was licensed to Pharmos for development from
 its originator, the Hebrew University of Jerusalem. Pharmos is seeking to
 enter into a strategic agreement with another company to develop and
 commercialize dexanabinol. Unlike its enantiomer, HU-210 (Yisum Research
 Development Co), dexanabinol does not interact with **cannabinoid**
 receptors. It has also exhibited more effective antioxidant and
 anti-inflammatory properties than MK-801 (dizocilpine; Merck & Co Inc).
 In addition, dexanabinol is generally well tolerated and appears toxicol.
 safe. Pharmos has been awarded a Small Business Innovation Research grant
 from the National Institutes of Health (NIH) National Institute of Neurol.
 Disorders and Stroke, Division of Stroke and Trauma. The grant covers the
 development of new prodrugs and novel formulations of dexanabinol and will
 support addnl. study of dexanabinol compds. for various indications. The
 prodrugs being studied are part of the group of compds. that include
 dexanabinol. A Notice of Allowance was received in Mar. 1999 on a patent
 covering the use of the drug in the treatment of MS. The use of
 dexanabinol and its derivs. to treat MS is described in US-05932610. An
 oral formulation of dexanabinol is claimed in US-05891468. Dexanabinol
 analogs with special utility in acute and chronic pain are claimed in
 US-04876276, while dexanabinol analogs for neuroprotection are claimed in
 US-06096740. Pharmos ests. that the worldwide market for dexanabinol in
 the treatment of severe head trauma may reach \$1 billion per yr.

IT 112924-45-5P, Dexanabinol


RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or
 effector, except adverse); BPR (Biological process); BSU (Biological

study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses) (therapeutic uses of dexanabinol)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(therapeutic uses of dexanabinol)

REFERENCE COUNT: 106 THERE ARE 106 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 2 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2000:724276 HCAPLUS

DOCUMENT NUMBER: 134:25030

TITLE: Dexanabinol (HU-211): a **nonpsychotropic cannabinoid** with neuroprotective properties

AUTHOR(S): Shohami, Esther; Mechoulam, Raphael

CORPORATE SOURCE: Department of Pharmacology, School of Pharmacy, The Hebrew University, Jerusalem, 91120, Israel

SOURCE: Drug Development Research (2000), 50(3/4), 211-215

CODEN: DDREDK; ISSN: 0272-4391

PUBLISHER: Wiley-Liss, Inc.

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review with 34 refs. The synthetic **cannabinoid** 1',1'-dimethylheptyl-(+)-(6aS,10aS)-11-hydroxy- Δ^8 -**tetrahydrocannabinol** (dexanabinol, HU-211) is inactive as a **cannabimimetic**, but exhibits pharmacol. properties characteristic of an N-methyl-D-aspartate (NMDA)-receptor antagonist. It blocks NMDA receptors stereospecifically by interacting with a site close to, but distinct from, that of uncompetitive NMDA receptor antagonists and from the recognition sites of glutamate, glycine, and polyamines. HU-211 inhibits the synthesis of tumor necrosis factor- α (TNF α) and possesses antioxidant properties. HU-211 blocked NMDA-induced ^{45}Ca uptake by primary neuronal cultures of rat forebrain and protected the same neuronal cultures against NMDA and glutamate neurotoxicity. Moreover, HU-211 effectively scavenged peroxy radicals in vitro and protected cultured neurons from the toxic effects of reactive O species (ROS). In addition, HU-211 markedly suppressed in vitro TNF α production and NO

generation (by >90%) by both murine peritoneal macrophages and rat alveolar macrophages exposed to lipopolysaccharide. Since glutamate, ROS and TNF α are implicated in the pathophysiol. of various acute conditions, the promising results showing neuroprotection by HU-211, acting via multiple mechanisms, led to a series of studies in which the drug was given to exptl. animals. This review discusses results from expts. describing the potential use of HU-211 as a neuroprotective agent in models of traumatic brain injury, stroke, optic nerve injury, pneumococcal meningitis, sepsis, and soman toxicity. In addition, HU-211 has been introduced into clin. trials for traumatic brain injury, and the successful results of two phases of clin. trials in head-injured patients are also shown.

IT 112924-45-5, Dexanabinol


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(dexanabinol (HU-211): a **nonpsychotropic cannabinoid** with neuroprotective properties)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 3 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2000:614231 HCAPLUS

DOCUMENT NUMBER: 133:275752

TITLE: **Nonpsychotropic synthetic cannabinoids**

AUTHOR(S): Pop, Emil

CORPORATE SOURCE: Alchem Laboratories Corporation, Alachua, FL, USA

SOURCE: Current Pharmaceutical Design (2000), 6(13), 1347-1359

CODEN: CPDEFP; ISSN: 1381-6128

PUBLISHER: Bentham Science Publishers

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review with 61 refs. Unlike natural **cannabinoids** which belong to the 6aR - trans series, the synthetic dexanabinol (HU-211), a 6aS-trans enantiomer, does not have affinity toward **cannabinoid** receptors and is devoid of **cannabimimetic** activity. On the other hand, dexanabinol demonstrated significant neuroprotective properties which prompted its development as a therapeutic agent. We now present the

extension of a series of 6aS-trans **cannabinoids** with novel derivs., including water soluble derivs. and congeners of dexanabinol.


IT 112924-45-5, Dexanabinol

RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(nonpsychotropic synthetic **cannabinoids**)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 4 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2000:614228 HCAPLUS

DOCUMENT NUMBER: 133:260976

TITLE: Looking back at **cannabis** research

AUTHOR(S): Mechoulam, Raphael

CORPORATE SOURCE: Medical Faculty, Department of Medicinal Chemistry and Natural Products, Hebrew University, Jerusalem, 91120, Israel

SOURCE: Current Pharmaceutical Design (2000), 6(13), 1313-1322

CODEN: CPDEFP; ISSN: 1381-6128

PUBLISHER: Bentham Science Publishers

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review with 70 refs. Research leading to the isolation of the plant **cannabinoids** during the 1960's and to the endogenous **cannabinoids**, during the 1990's is described. Investigations on two **non-psychotropic**, synthetic **cannabinoids**, HU-211, a neuroprotective agent and HU-308, a specific CB2 agonist are presented.


IT 112924-45-5, HU-211

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
(looking back at **cannabis** research)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 70 THERE ARE 70 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 5 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2000:430923 HCAPLUS

DOCUMENT NUMBER: 133:290851

TITLE: A submicron emulsion of HU-211, a synthetic **cannabinoid**, reduces intraocular pressure in rabbits

AUTHOR(S): Naveh, Nava; Weissman, Channa; Muchtar, Shaul; Benita, Shimon; Mechoulam, Raphael

CORPORATE SOURCE: Glaucoma and Pharmacology Laboratory, Goldschleger Eye Research Institute, Tel Hashomer, 52621, Israel

SOURCE: Graefe's Archive for Clinical and Experimental

Ophthalmology (2000), 238(4), 334-338

CODEN: GACODL; ISSN: 0721-832X

PUBLISHER: Springer-Verlag

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Purpose: To study the ocular hypotensive effect of a **nonpsychotropic cannabinoid**, HU-211 (11-hydroxy-Δ8-tetrahydrocannabinol, dimethylheptyl), an N-methyl-D-aspartate (NMDA) agonist, in normotensive rabbits. Methods: The **cannabinoid** HU-211, being lipophilic, was incorporated into a stable oil-in-water submicron sterile emulsion, consisting of 0.12% (weight/weight) HU-211. A single-dose, randomized and double-masked study was designed, using a Digilab 30R pneumotonometer to measure intraocular pressure (IOP) in normotensive rabbits. Results: Application of a single dose of HU-211 ophthalmic preparation resulted in an IOP reduction of 5.3 mmHg (24% of baseline),

first evident at 1.5 h post application and persisting for over 6 h. A small but significant lowering of pressure (12.5% of baseline) occurred in the contralateral eyes of HU-211 treated rabbits, lasting for 4 h post treatment. Conclusion: Our work demonstrated that HU-211, incorporated into submicron emulsion, caused a 6-h-long reduction in IOP in the treated eye, with a lesser reduction in the contralateral untreated eye.

IT 112924-45-5, HU-211


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(a submicron emulsion of HU-211, a synthetic **cannabinoid**, reduces intraocular pressure in rabbits)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 6 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2000:379603 HCAPLUS

DOCUMENT NUMBER: 133:115071

TITLE: Pharmacology of the intraocular pressure (IOP) lowering effect of systemic dexanabinol (HU-211), a **non-psychotropic cannabinoid**

AUTHOR(S): Beilin, Mark; Neumann, Ron; Belkin, Michael; Green, Keith; Bar-Ilan, Amir

CORPORATE SOURCE: Pharmos Corporation, Ltd., Rehovot, Israel

SOURCE: Journal of Ocular Pharmacology and Therapeutics (2000), 16(3), 217-230
CODEN: JOPTFU; ISSN: 1080-7683

PUBLISHER: Mary Ann Liebert, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The purpose of this study was to characterize the intraocular pressure (IOP) lowering activity and possible mechanism of action of the synthetic, **non-psychotropic cannabinoid** dexanabinol (HU-211) [(+)(3S,4S), 7-hydroxy- Δ -6- **tetrahydrocannabinol** 1, 1 dimethylheptyl], following i.v. administration in the rabbit. IOP (pneumatometry), aqueous humor inflow rate (fluorophotometry), blood pressure, and heart rate (computerized physiograph system connected to central ear artery cannula) were measured in unanesthetized albino rabbits. I.v. administration of HU-211 resulted in a dose-related reduction in IOP; a maximal IOP reduction of 5.0 ± 0.2 mmHg was observed 4 h after a 0.5 mg/kg dose. No significant changes in blood pressure or heart rate were observed during the first hr following this dose of HU-211. Pupil diameter did not change significantly during the 5 h following the 0.5 mg/kg i.v. dose. No significant change in the rate of aqueous humor inflow occurred during the 6 h after a 0.5 mg/kg dose of HU-211, thereby implicating outflow changes as the major source of IOP reduction. IOP reduction by HU-211 following pre-treatment with the α 2 adrenergic antagonist, yohimbine (1 mg/kg, i.v.), was only 30% of that of HU-211 alone. IOP reduction following pretreatment with the α 2 agonist, clonidine (0.5 mg/kg i.v.), was twice as large as that of HU-211 alone. Pretreatment with the β -adrenergic antagonist, propranolol (0.5 mg/kg i.v.), resulted in a 50% reduction in the IOP-lowering effect of HU-211. In summary, HU-211, administered i.v., is an effective IOP-lowering agent, devoid of any significant side effects (blood pressure, heart rate or pupil diameter, all of which have been reported previously for **cannabinoids**). Involvement of the adrenergic system is indicated in mediating the

IOP-lowering effects of HU-211 that appear to reflect a change in fluid outflow from the eye.

IT 112924-45-5, HU-211


RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(pharmacol. of intraocular pressure lowering effect of dexamabinol)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 7 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2000:74536 HCAPLUS

DOCUMENT NUMBER: 132:344219

TITLE: Neuroprotective effects of HU-211 on brain damage resulting from soman-induced seizures

AUTHOR(S): Filbert, Margaret G.; Forster, Jeffrey S.; Smith, C. Dahlem; Ballough, Gerald P. H.

CORPORATE SOURCE: U.S. Army Medical Research Institute of Chemical Defense, Aberdeen Proving Ground, MD, 21010-5400, USA

SOURCE: Annals of the New York Academy of Sciences (1999), 890 (Neuroprotective Agents), 505-514
CODEN: ANYAA9; ISSN: 0077-8923

PUBLISHER: New York Academy of Sciences

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The neuroprotective effects of HU-211 (dexamabinol); a synthetic **nonpsychotropic** analog of **tetrahydrocannabinol**, on brain damage resulting from soman-induced seizures were examined in male Sprague-Dawley rats challenged with 1.6 LD50 soman. At 5 or 40 min after onset of seizures, the rats were given an i.p. injection of 25 mg HU-211/kg. All rats that received soman showed electrocorticog. evidence of sustained seizures and status epilepticus for 4-6 h. HU-211 had no effect on either the strength or duration of seizure activity. Administration of HU-211 5 min after seizure onset reduced median lesion volume 86% (as assessed by microtubule-associated protein 2-neg. staining), and when administered 40 min postonset, the reduction in necrosis was 81.5% despite the presence of continuous seizures for 4-5 h. These observations were corroborated by hemotoxylin and eosin histopathol. assessment that showed a significant reduction in piriform cortical neuronal damage in HU-211-treated animals. It is concluded that HU-211 provides considerable

neuroprotection against brain damage produced by soman-induced seizures.

IT 112924-45-5, HU 211


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(neuroprotective effects of HU-211 on brain damage from soman-induced seizures)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 8 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1999:806942 HCAPLUS

DOCUMENT NUMBER: 132:175690

TITLE: Dexanabinol (HU-211) effect on experimental autoimmune encephalomyelitis: implications for the treatment of acute relapses of multiple sclerosis

AUTHOR(S): Achiron, A.; Miron, S.; Lavie, V.; Margalit, R.; Biegon, A.

CORPORATE SOURCE: Multiple Sclerosis Center, Sheba Medical Center, Tel-Hashomer, Ramat Gan, 52621, Israel

SOURCE: Journal of Neuroimmunology (2000), 102(1), 26-31

CODEN: JNRIDW; ISSN: 0165-5728

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Dexanabinol (HU-211) is a synthetic **non-psychotropic cannabinoid** which suppresses TNF- α production in the brain and peripheral blood. The effects of dexanabinol in rat exptl. autoimmune encephalomyelitis (EAE) were studied using different doses, modes of administration and time regimes. Dexanabinol, 5 mg/kg i.v. given once after disease onset (day 10), significantly reduced maximal EAE score. Increasing the dose or treatment duration resulted in further suppression of EAE. Drug administration at earlier phases during disease induction was not effective. Histol. studies supported the clin. findings demonstrating reduction in the inflammatory response in the brain and spinal cord in animals treated with dexanabinol. The results suggest that dexanabinol may provide an alternative mode of treatment for acute exacerbations of multiple sclerosis (MS).

IT 112924-45-5, HU-211


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(dexanabinol (HU-211) effect on exptl. autoimmune encephalomyelitis: implications for treatment of acute relapses of multiple sclerosis)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 9 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1999:796665 HCAPLUS

DOCUMENT NUMBER: 132:132238

TITLE: HU-308: a specific agonist for CB2, a peripheral **cannabinoid** receptor

AUTHOR(S): Hanus, L.; Breuer, A.; Tchilibon, S.; Shiloah, S.; Goldenberg, D.; Horowitz, M.; Pertwee, R. G.; Ross, R. A.; Mechoulam, R.; Fride, E.

CORPORATE SOURCE: Department of Medicinal Chemistry and Natural Products, Medical Faculty, Hebrew University, Jerusalem, 91120, Israel

SOURCE: Proceedings of the National Academy of Sciences of the United States of America (1999), 96(25), 14228-14233

CODEN: PNASA6; ISSN: 0027-8424

PUBLISHER: National Academy of Sciences

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Two **cannabinoid** receptors have been identified: CB1, present in the central nervous system (CNS) and to a lesser extent in other tissues, and CB2, present outside the CNS, in peripheral organs. There is evidence for the presence of CB2-like receptors in peripheral nerve terminals. We report now that we have synthesized a CB2-specific agonist, code-named HU-308. This **cannabinoid** does not bind to CB1 ($K_i > 10 \mu\text{M}$), but does so efficiently to CB2 ($K_i = 22.7 \pm 3.9 \text{ nM}$); it inhibits forskolin-stimulated cAMP production in CB2-transfected cells, but does so much less in CB1-transfected cells. HU-308 shows no activity in mice in a tetrad of behavioral tests, which together have been shown to be specific for **tetrahydrocannabinol** (THC)-type activity in the CNS mediated by CB1. However, HU-308 reduces blood pressure, blocks defecation, and elicits anti-inflammatory and peripheral analgesic activity. The hypotension, the inhibition of defecation, the anti-inflammatory and

peripheral analgesic effects produced by HU-308 are blocked (or partially blocked) by the CB2 antagonist SR-144528, but not by the CB1 antagonist SR-141716A. These results demonstrate the feasibility of discovering novel **nonpsychotropic cannabinoids** that may lead to new therapies for hypertension, inflammation, and pain.


IT 1972-08-3, **Tetrahydrocannabinol**

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(HU-308: a specific agonist for CB2, a peripheral **cannabinoid** receptor)

RN 1972-08-3 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 10 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1998:708291 HCAPLUS

DOCUMENT NUMBER: 130:133827

TITLE: **Δ⁹ Tetrahydrocannabinol** and **cannabidiol** alter cytokine production by human immune cells

AUTHOR(S): Srivastava, Maya D.; Srivastava, B. I. S.; Brouhard, B.

CORPORATE SOURCE: Department of Pediatrics, Cleveland Clinic Foundation, Cleveland, OH, 44195, USA

SOURCE: Immunopharmacology (1998), 40(3), 179-185
CODEN: IMMUDP; ISSN: 0162-3109

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Marijuana, a widely abused drug in the US, and its derivs. (**cannabinoids**) have been used in AIDS and cancer patients for treatment of intractable nausea and cachexia. Yet, objective investigations of the effect of **cannabinoids** on the human immune system are few. The authors investigated the effect of **Δ⁹ tetrahydrocannabinol** (THC) and **cannabidiol** (CBD) on cytokine production in vitro by human leukemic T, B, eosinophilic and CD8+ NK cell lines as models. THC decreased constitutive production of IL-8, MIP-1α, MIP-1β, and RANTES and phorbol ester stimulated production of TNF-α, GM-CSF and IFN-γ by NK cells. It inhibited MIP-1β in HTLV-1 pos. B-cells but tripled IL-8, MIP-1α and MIP-1β in B-cells and MIP-1β in eosinophilic cells but doubled IL-8. Both **cannabinoids** strongly inhibited IL-10 production by HUT-78 T-cells. Results indicate that THC and **nonpsychotropic** CBD have complex lineage and derivative specific effects on cytokines

consistent with previous animal studies. These effects while of potential benefits in some inflammatory/autoimmune diseases may worsen HIV infection, tumorigenesis and allergic inflammation in the lung.

IT 1972-08-3, Δ^9 - **Tetrahydrocannabinol**

13956-29-1, **Cannabidiol**


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(Δ^9 - **tetrahydrocannabinol** and **cannabidiol**
alter cytokine production by human immune cells)

RN 1972-08-3 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 13956-29-1 HCAPLUS

CN 1,3-Benzenediol, 2-[(1R,6R)-3-methyl-6-(1-methylethenyl)-2-cyclohexen-1-yl]-5-pentyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 11 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1998:460663 HCAPLUS

DOCUMENT NUMBER: 129:184186

TITLE: **Cannabidiol** and (-)- Δ^9 -**tetrahydrocannabinol** are neuroprotective antioxidants

AUTHOR(S): Hampson, A. J.; Grimaldi, M.; Axelrod, J.; Wink, D.
CORPORATE SOURCE: Laboratory of Cellular and Molecular Regulation,
National Institutes of Mental Health, Bethesda, MD,
20892, USA

SOURCE: Proceedings of the National Academy of Sciences of the United States of America (1998), 95(14), 8268-8273
 CODEN: PNASA6; ISSN: 0027-8424
 PUBLISHER: National Academy of Sciences
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The neuroprotective actions of **cannabidiol** and other **cannabinoids** were examined in rat cortical neuron cultures exposed to toxic levels of the excitatory neurotransmitter glutamate. Glutamate toxicity was reduced by both **cannabidiol**, a **nonpsychoactive** constituent of marijuana, and the **psychotropic cannabinoid** (-) Δ^9 -**tetrahydrocannabinol** (THC). **Cannabinoids** protected equally well against neurotoxicity mediated by N-methyl-D-aspartate receptors, 2-amino-3-(4-butyl-3-hydroxyisoxazol-5-yl)propionic acid receptors, or kainate receptors. N-methyl-D-aspartate receptor-induced toxicity has been shown to be calcium dependent; this study demonstrates that 2-amino-3-(4-butyl-3-hydroxyisoxazol-5-yl)propionic acid/kainate receptor-type neurotoxicity is also calcium-dependent, partly mediated by voltage sensitive calcium channels. The neuroprotection observed with **cannabidiol** and THC was unaffected by **cannabinoid** receptor antagonist, indicating it to be **cannabinoid** receptor independent. Previous studies have shown that glutamate toxicity may be prevented by antioxidants. **Cannabidiol**, THC and several synthetic **cannabinoids** all were demonstrated to be antioxidants by cyclic voltammetry. **Cannabidiol** and THC also were shown to prevent hydroperoxide-induced oxidative damage as well as or better than other antioxidants in a chemical (Fenton reaction) system and neuronal cultures. **Cannabidiol** was more protective against glutamate neurotoxicity than either ascorbate or α -tocopherol, indicating it to be a potent antioxidant. These data also suggest that the naturally occurring, **nonpsychotropic cannabinoid**, **cannabidiol**, may be a potentially useful therapeutic agent for the treatment of oxidative neurol. disorders such as cerebral ischemia.


IT 1972-08-3, (-)- Δ^9 - **Tetrahydrocannabinol**
 13956-29-1, **Cannabidiol**

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (cannabidiol and (-)- Δ^9 - **tetrahydrocannabinol**
 as neuroprotective antioxidants)

RN 1972-08-3 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 13956-29-1 HCAPLUS

CN 1,3-Benzenediol, 2-[(1R,6R)-3-methyl-6-(1-methylethenyl)-2-cyclohexen-1-yl]-5-pentyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 12 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1997:749546 HCAPLUS

DOCUMENT NUMBER: 128:70533

TITLE: Protection against septic shock and suppression of tumor necrosis factor α and nitric oxide production by dexamabinol (HU-211), a **nonpsychotropic cannabinoid**

AUTHOR(S): Gallily, Ruth; Yamin, Aviva; Waksman, Yaakov; Ovadia, Haim; Weidenfeld, Joseph; Bar-Joseph, Avi; Biegon, Anat; Mechoulam, Raphael; Shohami, Esther

CORPORATE SOURCE: Department of Immunology, Faculty of Medicine, The Hebrew University, Jerusalem, Israel

SOURCE: Journal of Pharmacology and Experimental Therapeutics (1997), 283(2), 918-924

CODEN: JPETAB; ISSN: 0022-3565

PUBLISHER: Williams & Wilkins

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Dexamabinol, HU-211, a synthetic **cannabinoid** devoid of **psychotropic** effects, improves neurol. outcome in models of brain trauma, ischemia and meningitis. Recently, HU-211 was found to inhibit brain tumor necrosis factor (TNF α) production after head injury. In the present study, we demonstrate the ability of HU-211 to suppress TNF α production and to rescue mice and rats from endotoxic shock after LPS (Escherichia coli 055:B5) inoculation. In BALB/c mice, a dose of 10 mg/kg LPS, injected i.p., caused 57% and 100% mortality, at 24 and 48 h, resp. HU-211, administered i.p. 30 min before lipopolysaccharide (LPS), reduced lethality to 9 and 67% at these time points (P < .05). When coinjected with D-galactoseamine (i.p.), LPS was 100% lethal within 24 h, whereas eight hourly injections of HU-211 caused mortality of C57BL/6 mice to drop to 10% (P < .001). Administration of LPS to Sprague-Dawley rats resulted in a 30% reduction in the mean arterial blood pressure within 30 min, which persisted for 3 h. HU-211, given 2 to 3 min before LPS, completely abolished the typical hypotensive response. Furthermore, the drug also markedly suppressed in vitro TNF α production and nitric oxide generation (by >90%) by both murine peritoneal macrophages and rat alveolar macrophage cell line exposed to LPS. HU-211 may, therefore, have therapeutic implications in the treatment of TNF α -mediated

pathologies.

IT 112924-45-5, Dexanabinol


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(protection against septic shock and suppression of tumor necrosis factor α and nitric oxide production by dexanabinol)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 13 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1997:747199 HCAPLUS

DOCUMENT NUMBER: 128:74980

TITLE: Dimerization of dexanabinol by hydrogen bonding accounts for its hydrophobic character

AUTHOR(S): Pop, Emil; Brewster, Marcus E.

CORPORATE SOURCE: Pharmos Corporation, Alachua, FL, 32615, USA

SOURCE: International Journal of Quantum Chemistry (1997), 65(6), 1057-1064
CODEN: IJQCB2; ISSN: 0020-7608

PUBLISHER: John Wiley & Sons, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Dexanabinol (I), a dihydroxylated synthetic **cannabinoid**, is a member of the **nonpsychotropic** (+)-3S,4S enantiomeric series. Exptl. evidence suggests that I might form aggregates (e.g., dimers) in which the 2 OH (a phenol and an allylic alc.) groups are involved in H bonding. The extremely low solubility of I in H₂O implies that this interaction may not involve solvent mols. A theor. study of this phenomenon in the framework of the PM3 mol. approximation is described. Simple mol. models (PhOH and 1-cyclohexene-1-methanol) were initially examined, followed by extension of the calcns. to I. I dimers resulting from H bonding are more stable than the isolated mols., with the differences attributed to H-bonding energies. The phenolic hydroxy group of 1 mol. forms an H bond with the allylic OH group of the 2nd mol. and vice versa, resulting in dimers containing 2 H bonds. The H bonds are more stable (6.14 kcal/mol) and the complex formed is more favored energetically when the phenol groups act as H-bond donors and the allylic OH groups as acceptors. These interactions are also energetically more favored than those between I and H₂O (3.70 kcal/mol). The I dimer showed a lower dipole moment than

the monomer (1.211 vs. 2.221 D) as well as a much larger log P (11.16 vs. 5.90), indicating strong hydrophobic character. The optimized structure shows that the OH groups involved in H bonds are oriented toward the interior of the dimers, while the lipophilic side chains are oriented toward the exterior. These properties of the dimer may explain the low water solubility of I.

IT 112924-45-5, Dexanabinol


RL: PRP (Properties)

(monomer and hydrogen-bonded dimer; dimerization of dexanabinol by hydrogen bonding in relation to hydrophobicity)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 14 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1997:709388 HCAPLUS

DOCUMENT NUMBER: 128:2351

TITLE: Oxidative stress in closed-head injury: Brain antioxidant capacity as an indicator of functional outcome

AUTHOR(S): Shohami, Esther; Beit-Yannai, Elie; Horowitz, Michal; Kohen, Ron

CORPORATE SOURCE: Departments of Pharmacology, Pharmaceuticals and Physiology, Schools of Pharmacy and Medicine, Hebrew University, Jerusalem, Israel

SOURCE: Journal of Cerebral Blood Flow and Metabolism (1997), 17(10), 1007-1019

CODEN: JCBMDN; ISSN: 0271-678X

PUBLISHER: Lippincott-Raven

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review, with 106 refs. It has been suggested that reactive oxygen species (ROS) play a role in the pathophysiol. of brain damage. A number of therapeutic approaches, based on scavenging these radicals, have been attempted both in exptl. models and in the clin. setting. In an exptl. rat and mouse model of closed-head injury (CHI), we have studied the total tissue nonenzymic antioxidant capacity to combat ROS. A major mechanism for neutralizing ROS uses endogenous low-mol. weight antioxidants (LMWA). This review deals with the source and nature of ROS in the brain, along with the endogenous defense mechanisms that fight ROS. Special emphasis is placed on LMWA such as ascorbate, urate, tocopherol, lipoic

acid, and histidine-related compds. A novel electrochem. method, using cyclic voltammetry for the determination of total tissue LMWA, is described.

The

temporal changes in brain LMWA after CHI, as part of the response of the tissue to high ROS levels, and the correlation between the ability of the brain to elevate LMWA and clin. outcome are addressed. We relate to the beneficial effects observed in heat-acclimated rats and the detrimental effects of injury found in apolipoprotein E-deficient mice. Finally, we summarize the effects of cerebroprotective pharmacol. agents including the iron chelator desferal, superoxide dismutase, a stable radical from the nitroxide family, and HU-211, a **nonpsychotropic cannabinoid** with antioxidant properties. We conclude that ROS play a key role in the pathophysiol. of brain injury, and that their neutralization by endogenous or exogenous antioxidants has a protective effect. It is suggested, therefore, that the brain responds to ROS by increasing LMWA, and that the degree of this response is correlated with clin. recovery. The greater the response, the more favorable the outcome.

IT 112924-45-5, HU-211


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(oxidative stress and brain antioxidant capacity as indicator of functional outcome in closed-head injury)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 106 THERE ARE 106 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L56 ANSWER 15 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1997:610131 HCAPLUS

DOCUMENT NUMBER: 127:242793

TITLE: Clinical pharmacokinetics of escalating i.v. doses of dexamabinol (HU-211), a neuroprotectant agent, in normal volunteers

AUTHOR(S): Brewster, M. E.; Pop, E.; Foltz, R. L.; Reuschel, S.; Griffith, W.; Amselem, S.; Biegon, A.

CORPORATE SOURCE: Pharmos Corporation, Alachua, FL, USA


SOURCE: International Journal of Clinical Pharmacology and Therapeutics (1997), 35(9), 361-365

CODEN: ICTHEK; ISSN: 0946-1965

PUBLISHER: Dustri-Verlag Dr. Karl Feistle

DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The pharmacokinetics of dexanabinol (HU-211), a synthetic, **nonpsychotropic cannabinoid** with neuroprotectant action, was evaluated administering i.v. doses of 48, 100, and 200 mg. All administrations were well tolerated. The elimination of HU-211 was best fitted to a 3-compartment model with a rapid distribution half-life (<5 min), an intermediate phase half-life of approx. 90 min, and a slow terminal elimination half-life (9 h). The pharmacokinetics were linear over the evaluated dose range. The plasma clearance was high (1.7 L/min), and the volume of distribution was approx. 15 L/kg.
 IT **112924-45-5**, Dexanabinol
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
 (HU-211; clin. pharmacokinetics of escalating i.v. doses)
 RN 112924-45-5 HCAPLUS
 CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 16 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1996:466532 HCAPLUS
 DOCUMENT NUMBER: 125:168374
 TITLE: Neuroprotective (+)-3S,4S-**cannabinoids** with modified 5'-side chain
 AUTHOR(S): Pop, Emil; Browne, Clinton E.; Nadler, Varda; Biegon, Anat; Brewster, Marcus E.
 CORPORATE SOURCE: Pharmos Corporation, Alachua, FL, 32615, USA
 SOURCE: Bioorganic & Medicinal Chemistry Letters (1996), 6(13), 1553-1558
 CODEN: BMCLE8; ISSN: 0960-894X
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 125:168374
 AB The synthesis and evaluation of two novel **cannabinoids** belonging to the (+)-3S,4S **nonpsychotropic** series are described. These derivs. bind to the NMDA receptor but have lower affinities than dexanabinol (HU-211), the series benchmark. The novel compds. protect neurons against NMDA-induced toxicity in cortical cell cultures and have lower toxicity to host neurons than dexanabinol.
 IT **180415-80-9P**
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological


study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation of neuroprotective **cannabinoids** with modified side chain as NMDA receptor inhibitors)

RN 180415-80-9 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylethyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



L56 ANSWER 17 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1996:183947 HCAPLUS

DOCUMENT NUMBER: 124:250801

TITLE: A novel **nonpsychotropic cannabinoid**, HU-211, in the treatment of experimental pneumococcal meningitis

AUTHOR(S): Bass, Roman; Engelhard, Dan; Trembovler, Victoria; Shohami, Esther

CORPORATE SOURCE: Departments Pharmacology and Pediatrics, Hebrew University, Jerusalem, 91120, Israel

SOURCE: Journal of Infectious Diseases (1996), 173(3), 735-8

CODEN: JIDIAQ; ISSN: 0022-1899

PUBLISHER: University of Chicago Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Typical features of pneumococcal meningitis have been demonstrated in rats inoculated with *Streptococcus pneumoniae*. HU-211, a novel noncompetitive N-methyl-D-aspartate antagonist recently demonstrated to inhibit tumor necrosis factor- α production under various conditions, improves recovery in some exptl. models of brain injury. The present study tested the efficacy of HU-211 in combination with antimicrobial therapy in reducing brain damage in exptl. pneumococcal meningitis. *S. pneumoniae*-infected rats were treated with saline alone, ceftriaxone alone, or with a combination of ceftriaxone and HU-211 18 h after inoculation of the bacteria. Brain edema and blood-brain barrier impairment 48 h after infection were significantly ($P < .05$) reduced in rats treated with ceftriaxone-HU-211 compared with rats in other treatment groups. The results suggest that HU-211 when given concomitantly with antibiotics attenuates brain damage in the rat model of pneumococcal meningitis.


IT 112924-45-5, HU-211

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (treatment of exptl. pneumococcal meningitis with **nonpsychotropic cannabinoid** HU-211)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 18 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1996:111999 HCAPLUS

DOCUMENT NUMBER: 124:220236

TITLE: Derivatives of dexanabinol. I. Water-soluble salts of glycinate esters

AUTHOR(S): Pop, Emil; Liu, Zong Zheng; Brewster, Marcus E.; Barenholz, Yechezkel; Korablyov, Veronica; Mechoulam, Raphael; Nadler, Varda; Biegon, Anat

CORPORATE SOURCE: Pharmos Corporation, Alachua, FL, 32615, USA

SOURCE: Pharmaceutical Research (1996), 13(1), 62-9

CODEN: PHREEB; ISSN: 0724-8741

PUBLISHER: Plenum

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Glycinate ester-type water soluble derivs. of dexanabinol (HU-211) (I) a **non-psychotropic cannabinoid** with potential use in the treatment of brain damage were synthesized and evaluated as prodrugs or congeners. Conventional procedures were used for the synthesis of the novel derivs. Stability studies in water and blood (rat, dog, human) were performed by HPLC; NMDA receptor binding was determined by radio ligand [³H] MK-801-displacement; the neuroprotection and neurotoxicity studies were performed in cortical cell cultures. Glycinate, dimethylamine, diethylamine, trimethylammonium and triethylammonium acetates of I were synthesized. All compds. were relatively soluble and stable in water. The quaternary ammonium salt-type derivs. rapidly hydrolyzed to the parent drug in various types of blood including human. In vitro activity studies indicated that the novel derivs. possess NMDA receptor binding properties. The neuroprotecting properties manifested by some of the new derivs. were associated with very low neuronal cell toxicity and are credited to parent compound released by hydrolysis during the expts. rather than to intrinsic activity. The trimethylammonium and triethylammonium acetates of I are promising water-soluble prodrug candidates for I; the glycinate ester might be used as an active analog.

IT 112924-45-5, Dexanabinol

RL: BSU (Biological study, unclassified); MFM (Metabolic formation); PRP (Properties); RCT (Reactant); BIOL (Biological study); FORM (Formation, nonpreparative); RACT (Reactant or reagent)


(preparation of water-soluble salts of glycinate esters of dexanabinol as prodrugs with neuroprotectant and NMDA receptor-antagonist activity)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-

tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS) - (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 19 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1996:68712 HCAPLUS

DOCUMENT NUMBER: 124:165086

TITLE: HU-211, a novel noncompetitive N-methyl-D-aspartate antagonist, improves neurological deficit and reduces infarct volume after reversible focal cerebral ischemia in the rat

AUTHOR(S): Belayev, Ludmila; Busto, Raul; Zhao, Weizhao; Ginsberg, Myron D.

CORPORATE SOURCE: School Medicine, University Miami, Miami, FL, 33101, USA

SOURCE: Stroke (Dallas) (1995), 26(12), 2313-20
CODEN: SJCCA7; ISSN: 0039-2499

PUBLISHER: American Heart Association

DOCUMENT TYPE: Journal

LANGUAGE: English

AB HU-211 is a **nonpsychotropic cannabinoid** analog that has been shown to act as a functional N-methyl-D-aspartate receptor blocker. The authors investigated the neuroprotective efficacy of HU-211 in a model of reversible middle cerebral artery occlusion (MCAo) in rats. Male Wistar rats were anesthetized with halothane and subjected to 90 min of temporary MCAo by retrograde insertion of an intraluminal nylon suture, coated with poly-L-lysine, through the external carotid artery into the internal carotid artery and MCAo. The drug (HU-211 in cosolvent, 4 mg/kg IV) or vehicle was administered in a blinded fashion 70 min after onset of MCAo. Behavioral tests were evaluated during occlusion (60 min) and for a 3-day period after MCAo. Three days after MCAo, brains were perfusion-fixed, and infarct vols. were determined. HU-211 significantly improved the neurol. score compared with vehicle during the 3 days after MCAo. Treatment with HU-211 also significantly reduced both infarct volume (66.6 vs. 149.8 mm³) and brain swelling (2.61% vs. 6.66%) compared with vehicle-treated rats (in each group). These results demonstrate the neuroprotective ability of HU-211 in focal cerebral ischemia as judged by neurol. score, infarct size, and brain swelling. Reversible MCAo with the use of a poly-L-lysine-coated intraluminal suture proved to be a reliable and effective modification of this technique, yielding consistent results.

IT 112924-45-5, HU-211

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(novel noncompetitive methyl-D-aspartate antagonist HU-211 improves

neurol. deficit and reduces infarct volume after reversible focal cerebral ischemia in rat induced with poly-L-lysine intraluminal suture)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI). (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 20 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1995:988808 HCAPLUS

DOCUMENT NUMBER: 124:76318

TITLE: Post-ischemic administration of HU-211, a novel non-competitive NMDA antagonist, protects against blood-brain barrier disruption in photochemical cortical infarction in rats: a quantitative study
AUTHOR(S): Belayev, Ludmila; Busto, Raul; Watson, Brant D.; Ginsberg, Myron D.

CORPORATE SOURCE: Cerebral Vascular Disease Research Center, Department of Neurology (D4-5), University of Miami School of Medicine, PO Box 016960, Miami, FL, 33101, USA

SOURCE: Brain Research (1995), 702(1,2), 266-70

CODEN: BRREAP; ISSN: 0006-8993

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

AB We examined the effect of HU-211, a synthetic **non-psychotropic cannabinoid** with non-competitive N-methyl-D-aspartate (NMDA) antagonist properties, on blood-brain barrier (BBB) integrity after photochem. induced cortical infarction. Evans blue dye was used as a BBB permeability indicator after unilateral thrombotic cortical infarction was produced photochem. by 560 nm light irradiation of the cortex in male Wistar rats receiving rose bengal i.v. HU-211 was injected in a dose of 4 mg/kg i.v. 30 min after stroke. Fluorometric measurement of Evans blue was performed 24 h later in six brain regions. Treatment with HU-211 significantly decreased extravasation of dye into the area of infarct (406 ± 19 vs. 539 ± 33 $\mu\text{g/g}$, mean \pm S.E.M.) as well as other sites of the affected hemisphere (866 ± 68 vs. 1096 ± 68 $\mu\text{g/g}$) compared to the vehicle group. These data indicate that HU-211 is an effective drug in protecting against the effects of focal ischemia-induced BBB disruption in the rat and suggest that the drug may be an effective treatment against the ischemic cell death and BBB disruption that can occur clin. following a stroke or cardiac arrest.

IT 112924-45-5, HU-211


RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(NMDA antagonist HU-211 protection against ischemia-induced blood-brain barrier disruption)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 21 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1995:709346 HCAPLUS

DOCUMENT NUMBER: 123:102677

TITLE: HU-211, a **nonpsychotropic cannabinoid**, improves neurological signs and reduces brain damage after severe forebrain ischemia in rats

AUTHOR(S): Belayev, Ludmila; Bar-Joseph, Avi; Adamchik, Jana; Biegon, Anat

CORPORATE SOURCE: Department Pharmacology, Pharmos Corp., Rehovot, 76326, Israel

SOURCE: Molecular and Chemical Neuropathology (1995), 25(1), 19-33

CODEN: MCHNEM; ISSN: 1044-7393

PUBLISHER: Humana

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The purpose of the present study was to examine the dose-response relationship and the therapeutic time window for the synthetic **nonpsychotropic cannabinoid** (HU-211) as a neuroprotective agent in transient, severe forebrain ischemia in the rat. Adult Sprague-Dawley rats were subjected to 20 min common carotid artery occlusion (CCAO) 24 h after coagulation of both vertebral arteries. Thirty minutes after the onset of CCAO, rats received an i.v. injection of HU-211 2, 4, or 8 mg/kg in HPCD (n = 12, 18, and 11, resp.), or the appropriate vehicle (n = 20). Neurol. signs were scored daily for 3 d following ischemia. A significant improvement (p < 0.01, Kruskal-Wallis nonparametric test, followed by Mann-Whitney U-test, p < 0.05) of neurol. deficits by the 4 mg/kg dose of HU-211, was observed 24, 48, and 72 h after insult. On the third day post-CCAO, the rat brain was taken for histopathol. evaluation of the CA-1 sector of the hippocampus. Counts of viable neurons in the hippocampal CA1 field showed significantly more live cells in the HU-211 (4 mg/kg) treated animals (P < 0.001, ANOVA followed by Duncan's post-hoc test, p < 0.05). The drug was equally effective when given 30 and 60 min after ischemia, but neuroprotection was no longer significant after 3 h. We suggest that HU-211 may be a potential treatment for postischemic brain damage in human beings.

IT 112924-45-5, HU-211


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(nonpsychotropic cannabinoid HU-211 improves
neurol. signs and reduces brain damage after forebrain ischemia)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 22 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1995:413714 HCAPLUS

DOCUMENT NUMBER: 122:205007

TITLE: Long-term effect of HU-211, a novel non-competitive NMDA antagonist, on motor and memory functions after closed head injury in the rat

AUTHOR(S) : Shohami, Esther; Novikov, Mark; Bass, Roman

CORPORATE SOURCE: Department of Pharmacology, The Hebrew University
School of Pharmacy, Jerusalem, Israel

SOURCE: Brain Research (1995), 674(1), 55-62

CODEN: BRREAP; ISSN: 0006-8993

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

AB HU-211 is a synthetic, **non-psychotropic**

cannabinoid which acts as a non-competitive.NMDA antagonist and antioxidant. We studied the drug's therapeutic window as well as its long-term effect on cognitive and motor functions in a model of closed head injury (CHI) in the rat. A weight-drop device was used to induce CHI in ether anesthetized male rats. HU-211 (5 mg/kg) was administered i.v. to the exptl. groups. For the therapeutic window study, drug was injected at 4 or 6 h after CHI. Edema (water content) and clin. status (neurol. severity score, NSS) were evaluated at 24 h. Reduction of edema was slight, whereas improvement of NSS was significant when the drug was administered at 4 or 6 h ($P = 0.0023$ and 0.059 , resp.). To determine the drug's long-term effect, it was administered 1 h after CHI and addnl. doses were later given. NSS was evaluated for a period of 30 d. A single dose of HU-211 given 1 h post-CHI improved the clin. outcome during the 30 d period ($P < 0.01$). Repetitive doses of HU-211 injected during the post traumatic period had similar effects. Cognitive functions were evaluated in the Morris water maze, with rats trained either before or after CHI. CHI resulted in a highly significant impairment of these abilities, whereas HU-211 treatment 1 h after CHI improved performance. Our results indicate

that HU-211 is a potent cerebroprotective agent, with a therapeutic window of about 4 h. The beneficial response obtained even after a single dose was long lasting, and ameliorated impairment of both motor and cognitive functions following CHI.


IT 112924-45-5, HU-211

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(long-term effect of NMDA antagonist HU-211 on motor and memory functions after closed head injury)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 23 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1995:390030 HCAPLUS

DOCUMENT NUMBER: 122:178176

TITLE: Neuroprotective activity of HU-211, a novel NMDA antagonist, in global ischemia in gerbils

AUTHOR(S): Bar-Joseph, Avi; Berkovitch, Yafit; Adamchik, Jana; Biegon, Anat

CORPORATE SOURCE: Pharmos Corp, Rehovot, 76326, Israel

SOURCE: Molecular and Chemical Neuropathology (1994), 23(2/3), 125-35

CODEN: MCHNEM; ISSN: 1044-7393

DOCUMENT TYPE: Journal

LANGUAGE: English

AB HU-211, a **nonpsychotropic cannabinoid** and a noncompetitive NMDA antagonist, was tested in a global ischemia model in the Mongolian gerbil. Male Mongolian gerbils underwent a 10-min bilateral common carotid artery occlusion. HU-211, administered i.v. at 4 mg/kg, 30 min postischemia, induced significant neuroprotection of the CA1 subfield of the hippocampus. A dose-response study demonstrated an inverted U curve in which the 4 mg/kg dose induced the best neuroprotection in the CA1 subfield of the hippocampus ($p < 0.05$ ANOVA followed by Duncan's post-hoc test). The therapeutic window was then investigated, and in another study, HU-211 4 mg/kg were administered i.v. at 30, 60, 120, and 180 min postinsult. A significant neuroprotection was detected at 30 and 60 min administration postinsult.


IT 112924-45-5, HU-211

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(neuroprotective activity of HU-211 in global ischemia)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 24 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1994:315619 HCAPLUS

DOCUMENT NUMBER: 120:315619

TITLE: HU-211, a **non-psychoactive cannabinoid**, rescues cortical neurons from excitatory amino acid toxicity in culture
 AUTHOR(S): Eshhar, Nomi; Striem, Sarina; Biegon, Anat
 CORPORATE SOURCE: Pharmacol. Dep., Pharmos Ltd., Rehovot, 76326, Israel
 SOURCE: NeuroReport (1993), 5(3), 237-40
 CODEN: NERPEZ; ISSN: 0959-4965

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The present study examined potential neuroprotective effects of HU-211, a synthetic **non-psychoactive cannabinoid** with non-competitive NMDA antagonist properties on neurons exposed to various excitotoxins in culture. HU-211 was found to protect neurons from NMDA and quisqualate-induced toxicity but not that produced by AMPA or kainate. NMDA-mediated neurotoxicity was blocked by HU-211 in a dose dependent manner with an $EC_{50} = 3.8 \pm 0.9 \mu M$. Radioligand binding studies have shown that HU-211 inhibits the binding of [3H]MK-801 to rat forebrain membranes ($KI = 11.0 \mu M \pm 1.323$) in a competitive manner, but was unable to displace [3H]kainate and [3H]AMPA binding. These data suggest that the neuroprotective activity of HU-211 is directly associated with the NMDA receptor channel and possibly with the quisqualate receptor of the metabotropic class. Thus, HU-211 appears to act as an NMDA open channel blocker and shows promise as a novel neuroprotectant for clin. use.

IT 112924-45-5, HU-211


RL: PRP (Properties)

(neuroprotectant activity and mechanism of)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 25 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1994:95460 HCAPLUS

DOCUMENT NUMBER: 120:95460

TITLE: The **non-psychotropic cannabinoid** (+)-(3S,4S)-7-hydroxy- Δ^6 -**tetrahydrocannabinol** 1,1-dimethylheptyl (HU-211) attenuates N-methyl-D-aspartate receptor-mediated neurotoxicity in primary cultures of rat forebrain

AUTHOR(S): Nadler, Varda; Mechoulam, Raphael; Sokolovsky, Mordechai

CORPORATE SOURCE: George S. Wise Fac. Life Sci., Tel Aviv Univ., Tel Aviv-Jaffa, 69978, Israel

SOURCE: Neuroscience Letters (1993), 162(1-2), 43-5
CODEN: NELED5; ISSN: 0304-3940

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The **non-psychotropic cannabinoid** (+)-(3S,4S)-7-hydroxy- Δ^6 - **tetrahydrocannabinol** 1,1-dimethylheptyl (HU-211), a stereoselective inhibitor of the N-methyl-D-aspartate (NMDA) receptor, protects primary cultures of rat forebrain against NMDA receptor-mediated neurotoxicity. Cell mortality produced by exposure for 10 min to NMDA or glutamate was reduced to approx. 18 or 27%, resp., by application of 50 μ M HU-211 for 10-15 min during or after exposure of cultures to excitatory amino acid. This effect of HU-211 was dependent on its concentration ($EC_{50} = 8.7 \pm 4 \mu$ M). HU-211 also reduced the toxicity induced by brief exposure (10 min) to kainate or quisqualate, though less effectively. HU-211 may therefore prove useful as a non-psychoactive drug that protects against neurotoxicity mediated by the NMDA receptor.

IT 112924-45-5, HU-211


RL: BIOL (Biological study)

(NMDA receptors neurotoxicity inhibition by, in forebrain)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 26 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1993:16190 HCAPLUS

DOCUMENT NUMBER: 118:16190

TITLE: Effect of synthetic enantiomeric **cannabinoids** on platelet aggregation

AUTHOR(S): Nathan, I.; Agam, G.; Mechoulam, R.; Dvilansky, A.; Livne, A. A.

CORPORATE SOURCE: Dep. Clin. Biochem., Ben Gurion Univ. Negev, Beer Sheva, 84105, Israel

SOURCE: Canadian Journal of Physiology and Pharmacology (1992), 70(10), 1305-8

CODEN: CJPPA3; ISSN: 0008-4212

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effect of a synthetic pair of enantiomeric **cannabinoids** on platelet function was evaluated. The **nonpsychotropic** enantiomer, the 1,1-dimethylheptyl homolog of (+)-(3S,4S)-7-hydroxy-Δ-6- **tetrahydrocannabinol** (HU-211), was found to be more active in inhibiting ADP-induced platelet aggregation than the highly **psychotropic** (-)-enantiomer (HU-210). The related (+)-(3R,3R) **cannabinoid**, HU-213, which lacks the 7-hydroxy moiety, exerted its inhibitory effect within a wider range of concns. The results indicate a differentiation between **psychotropic** activity and inhibition of platelet aggregation in the **cannabinoid** group of compds.


IT 112924-45-5, HU-211

RL: BIOL (Biological study)
(platelet aggregation inhibition and **psychotropic** activity of, structure in relation to)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 27 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1992:51323 HCAPLUS

DOCUMENT NUMBER: 116:51323

TITLE: Suppression of neuropathic pain behavior in rats by a **non-psychotropic** synthetic **cannabinoid** with NMDA receptor-blocking properties

AUTHOR(S): Zeltser, R.; Seltzer, Z.; Eisen, A.; Feigenbaum, J. J.; Mechoulam, R.

CORPORATE SOURCE: Dep. Oral Maxillofacial Surg., Hadassah Univ. Hosp., Jerusalem, Israel

SOURCE: Pain (1991), 47(1), 95-103
CODEN: PAINDB; ISSN: 0304-3959

DOCUMENT TYPE: Journal

LANGUAGE: English

AB HU211 is a novel synthetic derivative of tetrahydro-**cannabinol** (THC), the active marijuana ingredient. The stereochem. of HU211 is enantiomeric to that of THC. In contrast to THC, HU211 is not **psychotropic**. This agent exhibits other types of biol. activities; it is a non-competitive NMDA receptor blocker and has antinociceptive activity when injected with cupric chloride. This study examined its effects in autotomy, a behavioral model of neuropathic pain. Autotomy, a behavior of self-mutilation of denervated areas, was induced in Sabra rats by cutting the sciatic and saphenous nerves. Injections of HU211 (2.5 mg/kg) with cupric chloride (0.8 mg/kg) every 2nd day markedly suppressed autotomy during the injection period by delaying its average onset day and reducing the incidence of severe autotomy. Moreover, suppression of autotomy was retained in the postinjection period (for at least 30 days) but only when the drug was injected i.p. Lesser effects were achieved by s.c. injections. Cupric chloride or HU211 alone were ineffective. The general behavior and open field motor activity indicated that the effects of HU211 with Cu++ on autotomy were not due to sedation or ataxia but presumably due to antinociception mediated by NMDA receptor blockade.

IT 112924-45-5, HU211


RL: BIOL (Biological study)

(suppression of neuropathic pain behavior by, NMDA receptor block in)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 28 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1990:30609 HCAPLUS

DOCUMENT NUMBER: 112:30609

TITLE: **Nonpsychotropic cannabinoid** acts

AUTHOR(S): as a functional N-methyl-D-aspartate receptor blocker
Feigenbaum, Jeffery J.; Bergmann, Felix; Richmond,
Saul A.; Mechoulam, Raphael; Nadler, Varda; Kloog,

CORPORATE SOURCE: Fac. Med., Hebrew Univ., Jerusalem, 91120, Israel

SOURCE: Proceedings of the National Academy of Sciences of the
United States of America (1989), 86(23),
9584-7

CODEN: PNASA6; ISSN: 0027-8424

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Binding studies using the enantiomers of the synthetic **cannabinoid**
7-hydroxy- Δ^6 - **tetrahydrocannabinol** 1,1-dimethylheptyl
homolog in preps. of rat brain cortical membranes reveal that the
(+)-(3S,4S) enantiomer of this compound (HU-211) blocks N-methyl-D-aspartate
(NMDA) receptors in a stereospecific manner and that the interaction
occurs at binding sites distinct from those of other noncompetitive NMDA
antagonists or of glutamate and glycine. Moreover, HU-211 induces
stereotypy and locomotor hyperactivity in mice and tachycardia in rat,
effects typically caused by NMDA receptor antagonists. HU-211 is also a
potent blocker of NMDA-induced tremor, seizures, and lethality in mice.
This compound may therefore prove useful as a **nonpsychoactive** drug
that protects against NMDA-receptor-mediated neurotoxicity.


IT 112924-45-5, HU 211

RL: BIOL (Biological study)
(as methylaspartate receptor antagonist)

RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-
tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 29 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1990:525 HCAPLUS

DOCUMENT NUMBER: 112:525

TITLE: Inhibition of cisplatin-induced emesis in the pigeon by a **nonpsychotropic** synthetic **cannabinoid**

AUTHOR(S): Feigenbaum, Jeffery J.; Richmond, Saul A.; Weissman, Yoram; Mechoulam, Raphael

CORPORATE SOURCE: Med. Fac., Hebrew Univ., Jerusalem, Israel

SOURCE: European Journal of Pharmacology (1989), 169(1), 159-65

CODEN: EJPHAZ; ISSN: 0014-2999

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The (+) enantiomer of the synthetic **cannabinoid**, 7-hydroxy- Δ^6 - **tetrahydrocannabinol**, dimethylheptyl homolog (HU-211), possesses antiemetic efficacy in the pigeon. However, unlike all antiemetic **cannabinoids** tested in the past, it is devoid of **psychotropic (cannabinoid)** activity. The antiemetic activity of HU-211 was determined in pigeons given 10 mg/kg, i.v., cisplatin, a widely used antitumor agent, which is also a potent emesis producing agent at this dose. This activity was compared with that of Δ^1 - **tetrahydrocannabinol** (Δ^1 -THC). HU-211 pretreatment elicited a dose-related inhibition of cisplatin vomiting, with the optimal dose of HU-211 (2.5 mg/kg) inhibiting emesis by nearly 90%. Δ^1 -THC in doses up to 5 mg/kg caused only an insignificant reduction in vomiting. The activity was increased in the presence of CuCl₂ (0.8 mg/kg). The optimal dose of Δ^1 -THC (5.0 mg/kg) with CuCl₂ diminished the total amount of vomit expelled (up to 90%). However, it failed to inhibit emesis in 50% of all animals tested, did not affect the time of onset of emesis, and was highly **psychotropic**. The optimal dose of HU-211 (2.5 mg/kg) with CuCl₂ inhibited emesis by 97%, delayed the time on onset of emesis in the very few animals that did vomit, and was completely **nonpsychotropic**. The curve for the antiemetic effect of HU-211 was U-shaped over a narrow dose range. Thus, complete separation of **psychotropic** and antiemetic activities is possible in the **cannabinoid** series.

IT 1972-08-3, Δ^1 - **Tetrahydrocannabinol**

112924-45-5, HU 211

RL: BIOL (Biological study)


(vomiting inhibition by, **psychotropic** activity in relation to)

RN 1972-08-3 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-

, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 112924-45-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L56 ANSWER 30 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1986:102303 HCAPLUS

DOCUMENT NUMBER: 104:102303

TITLE: High-affinity **cannabinoid** binding sites in brain membranes labeled with [3H]-5'-trimethylammonium Δ 8- **tetrahydrocannabinol**

AUTHOR(S): Nye, Jeffrey S.; Seltzman, Herbert H.; Pitt, Colin G.; Snyder, Solomon H.

CORPORATE SOURCE: Sch. Med., Johns Hopkins Univ., Baltimore, MD, 21205, USA

SOURCE: Journal of Pharmacology and Experimental Therapeutics (1985), 234(3), 784-91
CODEN: JPETAB; ISSN: 0022-3565

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The binding of 3H-labeled 5'-trimethylammonium Δ 8- **tetrahydrocannabinol** (TMA) [99469-31-5], a pos. charged analog of Δ 8-THC modified on the 5' C (a portion of the mol. not important for its psychoactivity), to rat neuronal membranes was studied. Unlabeled TMA inhibits field-stimulated contractions of the guinea-pig ileum (IC50 = 1 μ M) in the same presynaptic manner as Δ 9-THC [3H]TMA binds saturably and reversibly to brain membranes with high affinity (Kp = 89 nM) to apparently 1 class of site (Hill coefficient, 1.1). Highest binding

site d. occurs in the brain, but several peripheral organs also display specific binding. Detergent solubilizes the sites without affecting their pharmacol. properties. Mol. sieve chromatog. reveals a bimodal peak of [3H]TMA binding activity of approx. 60,000 daltons apparent mol. weight Δ^9 -THC competitively inhibits [3H]TMA binding potently ($K_i = 27$ nM) and stereoselectively. For some **cannabinoids** potency in behavioral and physiol. tests parallels their affinity for the [3H]TMA binding site. However, several **nonpsychotropic cannabinoids** are active at the binding site.


IT 521-35-7 1972-08-3 5957-75-5
 13956-29-1 17766-02-8 21366-63-2
 25654-31-3 27262-32-4 28646-40-4
 32794-91-5 33029-18-4 34675-49-5
 34984-78-6 36557-05-8 36913-21-0
 39690-06-7 51895-51-3 53989-32-5
 56354-06-4 57030-51-0 58434-44-9

RL: PROC (Process)

(binding of, to brain membrane, pharmacol. activity in relation to)

RN 521-35-7 HCAPLUS


CN 6H-Dibenzo[b,d]pyran-1-ol, 6,6,9-trimethyl-3-pentyl- (7CI, 8CI, 9CI) (CA INDEX NAME)



RN 1972-08-3 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 5957-75-5 HCAPLUS


CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,10,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 13956-29-1 HCAPLUS
 CN 1,3-Benzenediol, 2-[(1R,6R)-3-methyl-6-(1-methylethenyl)-2-cyclohexen-1-yl]-5-pentyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




RN 17766-02-8 HCAPLUS
 CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




RN 21366-63-2 HCAPLUS
 CN 1H-4-Oxabenzof[cyclobut]inden-8-ol, 1a,2,3,3a,8b,8c-hexahydro-1,1,3a-trimethyl-6-pentyl-, (1aS,3aR,8bR,8cR)- (9CI) (CA INDEX NAME)




RN 25654-31-3 HCAPLUS
 CN 1,3-Benzenediol, 2-[(2E)-3,7-dimethyl-2,6-octadienyl]-5-pentyl- (9CI) (CA INDEX NAME)

Double bond geometry as shown.




RN 27262-32-4 HCAPLUS
 CN 6H-Dibenzo[b,d]pyran-1,8-diol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,8S,10aR)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 28646-40-4 HCAPLUS
 CN 6H-Dibenzo[b,d]pyran-9-methanol, 6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 32794-91-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,9,10,10a-hexahydro-6,6-dimethyl-9-methylene-3-pentyl-, trans- (8CI, 9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 33029-18-4 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,10,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aS,10aS)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 34675-49-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 6a,7,8,10a-tetrahydro-1-hydroxy-6,6-dimethyl-3-pentyl-, (6aR,10aR)-rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 34984-78-6 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1,8-diol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,8R,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 36557-05-8 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 6a,7,8,10a-tetrahydro-1-hydroxy-6,6-dimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 36913-21-0 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1,8-diol, 6a,7,8,10a-tetrahydro-9-(hydroxymethyl)-6,6-dimethyl-3-pentyl-, [6aR-(6aα,8β,10aβ)]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 39690-06-7 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-carboxylic acid, 6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 51895-51-3 HCAPLUS
 CN 6H-Dibenzo[b,d]pyran-1-ol, 6,6,9-trimethyl-3-pentyl-, acetate (9CI) (CA INDEX NAME)




RN 53989-32-5 HCAPLUS
 CN 6H-Dibenzo[b,d]pyran-9-carboxylic acid, 1-hydroxy-6,6-dimethyl-3-pentyl- (9CI) (CA INDEX NAME)



RN 56354-06-4 HCAPLUS
 CN 6H-Dibenzo[b,d]pyran-9-carboxylic acid, 6a,7,8,10a-tetrahydro-1-hydroxy-6,6-dimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).




RN 57030-51-0 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1,8-diol, 6a,7,8,10a-tetrahydro-9-(hydroxymethyl)-6,6-dimethyl-3-pentyl-, (6aR,8R,10aR)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 58434-44-9 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-3-propanol, α -ethyl-6a,7,8,10a-tetrahydro-1-hydroxy-6,6,9-trimethyl- (9CI) (CA INDEX NAME)

L56 ANSWER 31 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1986:45598 HCAPLUS

DOCUMENT NUMBER: 104:45598

TITLE: Labeling of a **cannabinoid** binding site in brain with a [3H]quaternary ammonium analog of delta-8-THC

AUTHOR(S): Nye, J. S.; Seltzman, H. H.; Pitt, C. G.; Snyder, S. H.

CORPORATE SOURCE: Sch. Med., Johns Hopkins Univ., Baltimore, MD, 21205, USA

SOURCE: Marihuana '84 [Eighty-Four], Proc. Oxford Symp.

Cannabis (1985), Meeting Date 1984, 253-62.


Editor(s): Harvey, D. J. IRL: Oxford, UK.

CODEN: 54PAAW

DOCUMENT TYPE: Conference

LANGUAGE: English

GI



AB The binding of ^3H -labeled 5'-trimethylammonium-8Δ-tetrahydrocannabinol (I) [99742-01-5] to rat neuronal membranes was studied. Unlabeled I inhibits field-stimulated contractions of the guinea pig ileum ($\text{IC}_{50} = 1000 \text{ nM}$) in the same presynaptic manner as 9Δ-THC. ^3H I binds saturably and reversibly to brain membranes with high affinity ($K_d = 89 \text{ nM}$) to apparently 1 class of sites (Hill coefficient = 1.1). The highest receptor site d. occurs into brain, but several peripheral organs also display specific binding. Detergent solubilizes the sites without affecting their pharmacol. properties. Mol. sieve chromatog. reveals a bimodal peak of ^3H I binding activity, mol. weight approx. 60,000 daltons. (-)-trans-9Δ-Tetrahydrocannabinol [1972-08-3] competitively inhibits ^3H I binding potently ($K_i = 27 \text{ nM}$) and stereoselectivity. For some cannabinoids potency in behavioral and physiol. tests parallels their affinity for the ^3H TMA binding site. However, several non-**psychotropic cannabinoids** are active at the binding site as well.


IT 521-35-7 1972-08-3 5957-75-5
 13956-29-1 17766-02-8 21366-63-2
 25654-31-3 27179-28-8 27262-32-4
 28646-40-4 30432-08-7 33029-18-4
 34984-78-6 36403-90-4 36557-05-8
 36913-21-0 39690-06-7 51895-51-3
 53989-32-5 56354-06-4 57030-51-0
 58434-44-9

RL: PROC (Process)

(binding of, to **cannabinoid** receptors of brain)

RN 521-35-7 HCAPLUS


CN 6H-Dibenzo[b,d]pyran-1-ol, 6,6,9-trimethyl-3-pentyl- (7CI, 8CI, 9CI) (CA INDEX NAME)



RN 1972-08-3 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 5957-75-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,10,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,10aR) - (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 13956-29-1 HCAPLUS

CN 1,3-Benzenediol, 2-[(1R,6R)-3-methyl-6-(1-methylethenyl)-2-cyclohexen-1-yl]-5-pentyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).




RN 17766-02-8 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aS,10aS) - (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




RN 21366-63-2 HCAPLUS
 CN 1H-4-Oxabenzo[f]cyclobut[cd]inden-8-ol, 1a,2,3,3a,8b,8c-hexahydro-1,1,3a-trimethyl-6-pentyl-, (1aS,3aR,8bR,8cR) - (9CI) (CA INDEX NAME)




RN 25654-31-3 HCAPLUS
 CN 1,3-Benzenediol, 2-[(2E)-3,7-dimethyl-2,6-octadienyl]-5-pentyl- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



RN 27179-28-8 HCAPLUS
 CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,9,10,10a-hexahydro-6,6-dimethyl-9-methylene-3-pentyl-, (6aR,10aR) - (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 27262-32-4 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1,8-diol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,8S,10aR) - (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 28646-40-4 HCAPLUS


CN 6H-Dibenzo[b,d]pyran-9-methanol, 6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-3-pentyl-, (6aR,10aR) - (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 30432-08-7 HCAPLUS


CN 6H-Dibenzo[b,d]pyran-9-methanol, 1-hydroxy-6,6-dimethyl-3-pentyl- (8CI, 9CI) (CA INDEX NAME)



RN 33029-18-4 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,10,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aS,10aS) - (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 34984-78-6 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1,8-diol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,8R,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 36403-90-4 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,9,10,10a-hexahydro-6,6,9-trimethyl-3-pentyl-, (6aR,9R,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 36557-05-8 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 6a,7,8,10a-tetrahydro-1-hydroxy-6,6-dimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 36913-21-0 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1,8-diol, 6a,7,8,10a-tetrahydro-9-(hydroxymethyl)-6,6-dimethyl-3-pentyl-, [6aR-(6a α ,8 β ,10a β)]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 39690-06-7 HCAPLUS


CN 6H-Dibenzo[b,d]pyran-9-carboxylic acid, 6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 51895-51-3 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6,6,9-trimethyl-3-pentyl-, acetate (9CI) (CA INDEX NAME)



RN 53989-32-5 HCAPLUS


CN 6H-Dibenzo[b,d]pyran-9-carboxylic acid, 1-hydroxy-6,6-dimethyl-3-pentyl-
(9CI) (CA INDEX NAME)



RN 56354-06-4 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-carboxylic acid, 6a,7,8,10a-tetrahydro-1-hydroxy-
6,6-dimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 57030-51-0 HCAPLUS


CN 6H-Dibenzo[b,d]pyran-1,8-diol, 6a,7,8,10a-tetrahydro-9-(hydroxymethyl)-6,6-
dimethyl-3-pentyl-, (6aR,8R,10aR)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 58434-44-9 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-3-propanol, α-ethyl-6a,7,8,10a-tetrahydro-1-hydroxy-6,6,9-trimethyl- (9CI) (CA INDEX NAME)



L56 ANSWER 32 OF 32 HCAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1978:608920 HCAPLUS

DOCUMENT NUMBER: 89:208920

TITLE: Molar volume relationships and the specific inhibition of a synaptosomal enzyme by psychoactive **cannabinoids**AUTHOR(S): Greenberg, Jeffrey H.; Mellors, Alan; McGowan, John C.
CORPORATE SOURCE: Guelph-Waterloo Cent. Grad. Work Chem., Univ. Guelph, Guelph, ON, Can.SOURCE: Journal of Medicinal Chemistry (1978),
21(12), 1208-12

CODEN: JMCMAR; ISSN: 0022-2623

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The ability of lipophilic compds. to inhibit the mouse-brain synaptosomal enzyme lysophosphatidylcholine acyltransferase [9027-64-9] was determined in vitro. Psychoactive **cannabinoids**, at lower levels than those predicted from their capacity to cause anesthesia, inhibited the enzyme, whereas **nonpsychoactive cannabinoids** did not show specific inhibition. M volume correlations were used to distinguish the nonspecific inhibition of the enzyme shown by all lipid substances from the specific inhibition shown by the psychoactive **cannabinoids**. The ability of the compds. to inhibit synaptosomal acyltransferase was compared with their ability to expand erythrocytic membranes.

IT 521-35-7 1972-08-3 5957-75-5


25654-31-3 36557-05-8

RL: BIOL (Biological study)

(synaptosomal lysophosphatidylcholine acyltransferase inhibition by)

RN 521-35-7 HCAPLUS


CN 6H-Dibenzo[b,d]pyran-1-ol, 6,6,9-trimethyl-3-pentyl- (7CI, 8CI, 9CI) (CA INDEX NAME)



RN 1972-08-3 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 5957-75-5 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-1-ol, 6a,7,10,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 25654-31-3 HCAPLUS

CN 1,3-Benzenediol, 2-[(2E)-3,7-dimethyl-2,6-octadienyl]-5-pentyl- (9CI) (CA INDEX NAME)


Double bond geometry as shown.




RN 36557-05-8 HCAPLUS

CN 6H-Dibenzo[b,d]pyran-9-methanol, 6a,7,8,10a-tetrahydro-1-hydroxy-6,6-dimethyl-3-pentyl-, (6aR,10aR)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



=> => d stat que 17
L5 STR



NODE ATTRIBUTES:
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 18

STEREO ATTRIBUTES: NONE
L7 0 SEA FILE=REGISTRY SSS FUL L5

100.0% PROCESSED 184 ITERATIONS
SEARCH TIME: 00.00.01

0 ANSWERS

=>